

(U)

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188PORT SECURITY CLASSIFICATION
(U)

MAR 14 1990

b. RESTRICTIVE MARKINGS
NASECURITY CLASSIFICATION AUTHORITY
NA

c. DISTRIBUTION/AVAILABILITY OF REPORT

CLASSIFICATION/DOWNGRADING SCHEDULE
NA

Distribution Unlimited

FORMING ORGANIZATION REPORT NUMBER(S)

Georgia State University

5. MONITORING ORGANIZATION REPORT NUMBER(S)

NA

NAME OF PERFORMING ORGANIZATION

Georgia State University

6b. OFFICE SYMBOL
(If applicable)
NA

7a. NAME OF MONITORING ORGANIZATION

Office of Naval Research

ADDRESS (City, State, and ZIP Code)

Georgia State University
Atlanta, GA 30303

7b. ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street
Arlington, VA 22217-5000NAME OF FUNDING/SPONSORING
ORGANIZATION
Office of Naval Research8b. OFFICE SYMBOL
(If applicable)
ONR

9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER

N00014-89-J-1640

8c. ADDRESS (City, State, and ZIP Code)

800 N. Quincy Street
Arlington, VA 22217-5000

10. SOURCE OF FUNDING NUMBERS

PROGRAM
ELEMENT NO.PROJECT
NO.TASK
NO.WORK UNIT
ACCESSION NO.

61153N

RR04108

4414013

11. TITLE (Include Security Classification)

(U) Neurochemical Control of Circadian Rhythms

12. PERSONAL AUTHOR(S)

Albers, H. Elliott

13a. TYPE OF REPORT

Annual

13b. TIME COVERED

FROM 3/31/89 TO 2/28/90

14. DATE OF REPORT (Year, Month, Day)

2/20/90

15. PAGE COUNT

13

16. SUPPLEMENTARY NOTATION

17. COSATI CODES

FIELD

GROUP

SUB-GROUP

08

18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)

peptides; hypothalamus; co-localization; light; gene
expression.

19. ABSTRACT (Continue on reverse if necessary and identify by block number)

The suprachiasmatic nucleus (SCN), which appears to act as a circadian clock, contains a large subpopulation of local circuit neurons in which vasoactive intestinal peptide (VIP) and peptide histidine isoleucine (PHI) are co-localized. We are continuing to investigate the hypothesis that VIP/PHI-containing neurons are essential for the synchronization of circadian rhythms with the day-night cycle. Using *in situ* hybridization, we have demonstrated that some, but not all SCN neurons that contain VIP/PHI mRNA also contain the mRNA encoding a third biologically active peptide, gastrin releasing peptide (GRP). Furthermore, comparison of the cellular levels of VIP/PHI and GRP mRNA revealed that these mRNAs have distinctly different 24 hr patterns within the SCN, (Continue)

20. DISTRIBUTION/AVAILABILITY OF ABSTRACT

☒ UNCLASSIFIED/UNLIMITED ☐ SAME AS RPT. ☐ DTIC USERS

21. ABSTRACT SECURITY CLASSIFICATION

(U)

22a. NAME OF RESPONSIBLE INDIVIDUAL

Dr. J.A. Majce

22b. TELEPHONE (Include Area Code)

202-696-4055

22c. OFFICE SYMBOL

ONR

(U)

SECURITY CLASSIFICATION OF THIS PAGE

In other experiments we examined whether VIP, PHI and GRP interact within the SCN to produce a signal important for circadian control. Co-administration of VIP, PHI and GRP within the SCN in vivo mimicked the phase delaying effects of light on circadian control, and activated SCN single units recorded in vitro. In contrast, administration of VIP, PHI or GRP alone, or co-administration of any two of these peptides did not produce the full behavioral or cellular response observed following co-administration of VIP, PHI and GRP. These data indicate that the interaction of VIP, PHI and GRP may be required for regulation of circadian rhythms by the SCN.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

INTRODUCTION

Several variations in the molecular mechanisms producing co-localized peptides have been observed. In some cases, the co-localized peptides come from the same precursor, while in other cases the co-localized peptides are derived from different precursors. One population of hypothalamic neurons in which both of these mechanisms appear to be active is in the suprachiasmatic nucleus (SCN). A large subpopulation of local circuit SCN neurons contain two peptides derived from the same precursor, VIP and peptide histidine isoleucine (PHI) and, also appear to contain a third peptide derived from a different precursor, gastrin releasing peptide (GRP) (Okamura, et. al., 1986).

A large body of evidence supports the view that the SCN acts as a circadian clock that both generates rhythms in behavior and physiology, and synchronizes those rhythms with the environmental light-dark (LD) cycle (Minors & Waterhouse, 1986). The SCN can be divided into anatomically distinct subpopulations of intrinsic neurons, however it is not known how these different neuronal subpopulations contribute to the generation and synchronization of circadian rhythms with the LD cycle (Moore & Card, 1985). Previously, we hypothesized that VIP/PHI-containing SCN neurons are involved in mediating the effects of environmental lighting important in the synchronization of circadian rhythms with the LD cycle. This hypothesis is supported by the findings that environmental lighting and enucleation selectively alter VIP and PHI immunoreactivity and mRNA within the SCN, and that the cellular levels of VIP/PHI mRNA vary over the day-night cycle (Albers, et. al., 1987; Stopa, et. al., 1988; Albers, et. al., 1990). The majority of VIP/PHI immunoreactive neurons are contained within the ventrolateral SCN, which is also the site of termination of most SCN afferent pathways, including two well defined photic projections, the retinohypothalamic tract (RHT) and the geniculohypothalamic tract (GHT). The possibility that VIP, PHI and GRP are co-localized in local circuit neurons in the afferent terminal fields of the SCN suggest that synchronization of circadian rhythms with the LD cycle may depend on the co-release of these three peptides. If so, the combined effect of VIP, PHI and GRP on circadian rhythms should be different from the effects of each peptide given alone, or in combination with one of the other two peptides. In the last year we have demonstrated that: 1) VIP/PHI and GRP mRNAs can be co-localized in SCN neurons, 2) VIP and GRP mRNAs exhibit different 24 hr patterns within the SCN, and 3) VIP, PHI and GRP interact to phase shift circadian activity rhythms and activate single unit discharge within the SCN.

METHODS

Microinjection Experiments

Hamsters were deeply anesthetized with sodium pentobarbital

and stereotaxically implanted with 26 gauge guide cannulas aimed at the suprachiasmatic region. The stereotaxic coordinates were 1.6 mm anterior and 1.7 mm lateral to bregma, and -7.7 mm below dura. The incisor bar was set at 0 and the stereotaxic arm was set 10 degrees from perpendicular. Peptides were dissolved in 0.9% NaCl (saline) and injected with a 1 μ l Hamilton syringe connected to a 33-gauge needle by polyethylene tubing. Peptides were purchased from Bachem, Inc, Torrance, CA. A cocktail containing approximately equimolar concentrations of VIP, PHI and GRP was made by dissolving approximately 50 pmol VIP, 50 pmol PHI and 50 pmol GRP into 400 μ l of saline. In all protocols a total of 150 pmol of peptide was injected in 400 μ l of saline. Following the experiment hamsters were deeply anesthetized with sodium pentobarbital and microinjected with 400 μ l of dye. Following intracardial perfusion with Perfix (Fisher Scientific), the brain was removed and examined to verify the site of injection. Phase shifts in the activity rhythm were calculated by linear regression.

Hypothalamic Slice Experiments

In brief, coronal brain slices containing the SCN (400-450 μ m thick) were harvested from hamsters housed in a LD 14:10 cycle. Immediately after sectioning, the slices were placed in incubation medium (i.e. modified Krebs solution) oxygenated with 95% O₂ and 5% CO₂ at 35 °C for at least one hr. Extracellular recordings were obtained using glass micropipettes with DC resistances of 25-40 M Ω filled with 0.5 M sodium acetate containing 2% Fast green. The micropipette was introduced into the slice with the aid of a microscope. At the end of the recording session, a current of 5 μ A was passed through the electrode for 3-5 min to deposit a blue spot. All recording sites used in the present study were in the ventrolateral SCN, where the VIP, PHI and GRP immunoreactive neurons are primarily located.

in situ hybridization

Rats maintained in a LD 14:10 cycle were sacrificed by decapitation at 4 hr intervals (N=4-7 per time point). Twelve micron thick coronal sections were cut through the anterior-posterior extent of the SCN. Hybridizations were performed using synthetic oligonucleotide probes specific to GRP and VIP mRNAs by methods described previously (Albers, et. al., 1990). The probes recognizing GRP and VIP mRNA were used for hybridizations conducted on adjacent coronal sections of the SCN. Cellular levels of the hybridization signal were determined with Bioquant Image Analysis software.

Statistical Analyses

Statistical differences among groups were assessed with the one-way classification of the analysis of variance. Subsequent to the analysis of variance, a priori comparisons were made using t-tests and post hoc comparisons between individual groups were made with Duncan's Multiple Range test.

RESULTS AND DISCUSSION

Co-localization of VIP/PHI and GRP mRNA within SCN Neurons

The co-localization of VIP/PHI and GRP in SCN neurons was examined by comparison of VIP/PHI and GRP mRNA within neurons of the ventrolateral SCN using in situ hybridization. As expected, the autoradiographic signal produced by probes complementary to VIP/PHI and GRP mRNAs was restricted to the ventrolateral aspect of the SCN. Analysis of adjacent sections labeled separately for VIP/PHI mRNA or GRP mRNA indicated that GRP mRNA-containing neurons also appeared to contain VIP/PHI mRNA. However, not all neurons displaying hybridization signal for VIP/PHI mRNA also contained hybridization signal for GRP. In summary, these studies suggest that some SCN neurons contain both VIP/PHI and GRP mRNA, while others contain only VIP/PHI mRNA.

Rhythms of VIP and GRP mRNA within the SCN

Comparison of the cellular levels of VIP/PHI and GRP mRNA in alternate sections of the SCN by in situ hybridization revealed that these mRNAs displayed distinctly different 24 hr patterns (Figure 1). As we had seen previously (Albers, et. al., 1990), VIP/PHI mRNA was greater at night than during the day. In contrast, GRP hybridization signal was greater during the day than at night. These data indicate that the ratio of VIP/PHI to GRP within the SCN may vary over the circadian cycle, and thereby could represent a "time of day" signal. In the upcoming year we will investigate whether the rhythmicity of afferent activity, or endogenous circadian activity induces the rhythms in VIP/PHI and GRP mRNA.

Behavioral Effects of VIP, PHI and GRP within the SCN

If VIP, PHI and GRP combine to form an important circadian signal within the SCN, co-administration of VIP, PHI and GRP within the SCN should influence the timing of circadian rhythms in vivo. To examine this possibility a cocktail containing equimolar concentrations of VIP, PHI and GRP was microinjected into the SCN region of hamsters maintained in constant illumination. As can be seen in Figure 2, VIP/PHI/GRP co-administered into the SCN region around the time of the onset of locomotor activity produced large phase delays in the free-running circadian activity rhythm. However microinjection of VIP/PHI/GRP at other times within the circadian cycle did not phase shift the activity rhythm (Figure 3). The phase shifts produced by VIP/PHI/GRP were the result of the action of these peptides in the SCN region, since microinjection of VIP/PHI/GRP into the cerebroventricular system did not phase shift the activity rhythm. Comparison of the phase response curve for VIP/PHI/GRP microinjection with the phase response curves for various lighting stimuli revealed that VIP/PHI/GRP mimicked the phase delaying, but not the phase advancing effects of light

pulses.

Although VIP/PHI/GRP mimicked the phase delaying effects of light within the SCN region, it was important to determine whether an interaction between all three peptides is required to produce these circadian effects, or whether the action of one or two of these peptides was sufficient. The phase shifting effects of administration of VIP, PHI and GRP and the co-administration of VIP/PHI, VIP/GRP and PHI/GRP were compared with the phase shifting effects of co-administration of VIP/PHI/GRP. The phase delays produced by microinjection of VIP/PHI/GRP were nearly two-fold greater than the phase delays produced by any of these peptides microinjected alone, or in combination with one of the others (Figure 4, Top).

Cellular Effects of VIP, PHI and GRP within the SCN

The hypothalamic slice preparation was used to investigate the cellular effects VIP, PHI and GRP on spontaneously active SCN neurons. Administration of a cocktail containing VIP/PHI/GRP into the perfusate was found to increase the spontaneous discharge of SCN neurons in a dose-dependent manner. At the phase of the circadian cycle in which microinjection of VIP/PHI/GRP phase delayed locomotor rhythms, the firing rate of neurons recorded in the ventrolateral SCN increased by approximately 4 impulses/s in response to VIP/PHI/GRP. The excitation produced by VIP/PHI/GRP was significantly larger than that produced by VIP, PHI, GRP, VIP/PHI, VIP/GRP or PHI/GRP (Figure 4, Bottom; Figure 5 and 6). These data indicate that VIP, PHI and GRP may function interactively within the SCN to regulate circadian rhythms.

References

Albers, H.E., Minamitani, N., Stopa, E.G. and Ferris, C.F. Light selectively alters vasoactive intestinal peptide and peptide histidine immunoreactivity within the rat suprachiasmatic nucleus. Brain Research, 437:189-192, 1987.

Albers, H.E., Stopa, E.G., Zoeller, R.T., Kauer, J.S., King, J.C., Fink, J.S., Mobtaker, H. and Wolfe, H. Day-night variation in prepro vasoactive intestinal peptide/peptide histidine isoleucine mRNA within the rat suprachiasmatic nucleus. Molecular Brain Research, 7:85-89, 1990.

Minors, D.S. and Waterhouse, J.M. Circadian rhythms and their mechanisms. Experientia, 42:1-13, 1986.

Moore, R.Y. and Card, J.P. Visual pathways and the entrainment of circadian rhythms. Annals New York Academy of Sciences, 453:123-133, 1985.

Okamura, H., Murakami, S., Uda, K., Sugano, T., Takahashi, Y., Yanaihara, C., Yanaihara, N. and Ibata, Y. Co-existence of vasoactive intestinal peptide (VIP)-, peptide histidine isoleucine amide (PHI)-, and gastrin releasing peptide (GRP)- like immunoreactivity in neurons of the rat suprachiasmatic nucleus. Biological Research, 7:295-299, 1986.

Stopa, E.G., Minamitani, N., Jonassen, J.A., King, J.C., Wolfe, H., Mobtaker, H. and Albers, H.E. Localization of vasoactive intestinal peptide and peptide histidine isoleucine immunoreactivity and mRNA within the rat suprachiasmatic nucleus. Molecular Brain Research, 4:319-325, 1988.

GRANT RELATED PUBLICATIONS

Abstracts

1. Albers, H.E., Liou, S.Y. and Zoeller, R.T. Peptide co-localization: Functional significance within the circadian timing system. Society for Neuroscience, Fall, 1989.
2. Albers, H.E., Liou, S.Y., Stopa, E.G. and Zoeller, R.T. Molecular, cellular and behavioral analysis of the circadian functions of vasoactive intestinal peptide (VIP), peptide histidine isoleucine (PHI) and gastrin releasing peptide (GRP). Symposium entitled Suprachiasmatic Nucleus: The Mind's Clock. National Institute of Child Health and Human Development, Fall, 1989.
3. Zoeller, R.T., Licklider, N.R., Anderson, E.R. and Albers, H.E. Cellular levels of messenger RNAs encoding VIP/PHI and gastrin releasing peptide (GRP, Bombesin) exhibit different 24 hr rhythms in the rat SCN. Endocrine Society, 1990, Submitted.

Papers

1. Albers, H.E., Stopa, E.G., Zoeller, R.T., Kauer, J.S., King, J.C., Fink, J.S., Mobtaker, H. and Wolfe, H. Day-night variation in vasoactive intestinal peptide/peptide histidine isoleucine mRNA within the rat suprachiasmatic nucleus. Molecular Brain Research, 7:85-89, 1990.
2. Albers, H.E., Ottenweller, J.E., Liou, S.Y., Lumpkin, M.D. and Anderson, E.R. Neuropeptide Y in the hypothalamus: Effect on corticosterone and single-unit activity. American Journal of Physiology, 258:R376-R382, 1990.
3. Albers, H.E., Liou, S.Y., Ferris, C.F., Stopa, E.G. and Zoeller, R.T. Neurochemistry of circadian timing. In: The Suprachiasmatic Nucleus: The Mind's Clock. D. Klein, R.Y. Moore, and S. Reppert (Eds.), Oxford University Press.
4. Liou, S.Y., Shibata, S., Albers, H.E. and Ueki, S. Effects of GABA and Anxiolytics on the single unit discharge of suprachiasmatic neurons in rat hypothalamus slices. Submitted.
5. Liou, S.Y. and Albers, H.E. Single unit response of neurons within the suprachiasmatic nucleus to GABA and

low chloride perfusate during day and night. Submitted.




Figure 1. Patterns of the relative cellular levels of GRP and VIP mRNA within the SCN of rats were housed in a LD cycle consisting of 14 hrs of light and 10 hrs of darkness. Both VIP and GRP mRNA were determined within the SCN of each rat by in situ hybridization. For both GRP and VIP mRNA each point is plotted as a percentage of the value observed at circadian time 4. However at all times of day the cellular levels of GRP mRNA were considerably lower than those of VIP mRNA.




Figure 2. Effect of various combinations of VIP, PHI and GRP on the phase of free-running activity rhythms following microinjection into the suprachiasmatic region of hamsters. Co-administration of VIP (50 pmol), PHI (50 pmol) and GRP (50 pmol) produced large phase delays when given around the time of activity onset, but had little effect at other times within the circadian cycle. Microinjection of VIP (150 pmol) or GRP (150 pmol), or co-administration of VIP (75 pmol) and GRP (75 pmol) or PHI (75 pmol) and GRP (75 pmol) produced only small delays in circadian phase.




Figure 3. Summary of the effects of microinjections of VIP/PHI/GRP (N=39) into the suprachiasmatic region plotted as a phase response curve. Solid circles represent mean phase shifts (\pm SEM), in hrs, at 3 hr intervals throughout the circadian cycle (circadian time 12 refers to the time of activity onset). VIP/PHI/GRP produced phase delay shifts around the time of activity onset, but had no effect at other times within the circadian cycle. VIP/PHI/GRP microinjection mimics the phase delaying but not the phase advancing effects of brief light pulses.




Figure 4. Summary of the behavioral and cellular effects of administration of VIP, PHI, GRP alone, co-administration of two of these peptides (i.e. VIP/PHI, VIP/GRP and PHI/GRP) and co-administration of all three peptides (VIP/PHI/GRP). TOP: Phase shifts produced in hamster activity rhythms following peptide microinjection into the suprachiasmatic region during the 3 hr interval following activity onset (i.e. circadian time 12-15). BOTTOM: Increase in spontaneous discharge of SCN single units following administration of VIP, PHI and GRP and co-administration of all possible combinations of these peptides in a concentration of 10^{-7} M. All single unit recordings were made during an 8 hr interval (circadian time 8-16) of the circadian cycle beginning 4 hrs before the time corresponding to activity onset (circadian time 12).




Figure 5. Integrated firing rate records from two suprachiasmatic neurons recorded extracellularly in the hypothalamic slice preparation. (A) Response of a SCN neuron to VIP, PHI, and co-administration of VIP/PHI. (B) Response of a SCN neuron to VIP, PHI, GRP and co-administration of VIP/PHI/GRP.




Figure 6. Dose-response of SCN single units to co-administration of VIP/PHI/GRP (N=5), VIP/PHI (N=4), VIP/GRP (N=3), PHI/GRP (N=3), VIP (N=5), PHI (N=5) and GRP (N=5).

DISTRIBUTION LIST

Stress Neurochemistry Program

Annual, Final and Technical Reports (one copy each)

INVESTIGATORS

Dr. H. Elliott Albers
Lab. Neuroendocrin. & Behavior
Depts. of Biology & Psychology
Georgia State University
Atlanta, GA 30303

Dr. Gwen V. Childs
Dept. of Anatomy & Neuroscience
Univ. of Texas Medical Branch
Galveston, TX 77550

Dr. Carl E. Creutz
Dept. of Pharmacology
University of Virginia
Charlottesville, VA 22908

Dr. Mary F. Dallman
Dept. of Physiology
University of California, Box 0444
San Francisco, CA 94143-0444

Dr. Caleb E. Finch
Dept. of Neurobiology
Univ. of Southern California
Los Angeles, CA 90089-0191

Dr. Thackery S. Gray
Department of Anatomy
Loyola University Medical Center
216 South First Avenue
Maywood, IL 60153

Dr. Richard F. Ochillo
College of Pharmacy
Xavier Univ. of Louisiana
7325 Palmetto Street
New Orleans, LA 70125

Dr. Terry Reisine
Dept. of Pharmacology
Univ. of Pennsylvania
School of Medicine
36th and Hamilton Walk
Philadelphia, PA 19104

Dr. C. Frank Starmer
P.O. Box 3181
Duke Univ. Medical Center
Durham, NC 27710

Dr. Kent E. Vrana
Dept. of Biochemistry
West Virginia School
of Medicine
Morgantown, WV 26506

Stress Neurochemisty

Annual, Final and Technical Reports (one copy each except as noted)

ADMINISTRATORS

Scientific Officer, Physiology
Code 1141SB
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

Program Manager, Code 1213
Human Factors Biosciences
Division
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217-5000

Administrator (2 copies) (Enclose DTIC Form 50)
Defense Technical Information Center
Building 5, Cameron Station
Alexandria, VA 22314

Program Manager, Code 223
Support Technology
Directorate
Office of Naval Technology
800 N. Quincy Street
Arlington, VA 22217-5000

Administrative Contracting Officer
ONR Resident Representative
(address varies - obtain from business office)

Annual and Final Reports Only (one copy each)

DoD ACTIVITIES

Commanding Officer
Naval Medical Center
Washington, DC 20372

Commanding Officer, Code 404
Naval Medical Research & Development Command
National Naval Medical Center
Bethesda, MD 20814

Commander
Chemical and Biological Sciences Division
Army Research Office, P.O. Box 12211
Research Triangle Park, NC 27709

Directorate of Life Sciences
Air Force Office of Scientific Research
Bolling Air Force Base
Washington, DC 20332

Final and Technical Reports Only

Director, Naval Research Laboratory (6 copies)
Attn: Technical Information Division, Code 2627
Washington, DC 20375